

ASTHMA HEALTH CARE PLAN

Child name:		Е	Birth date:		
Community program name:					
Parent/guardian name:					
Home Ph#:	Cell #:	V	Vork Ph#:		
Parent/guardian name:					
Home Ph#:	Cell #:		Work Ph#:		
Alternate emergency contact name	:				
Home Ph#:	Cell #:		Work Ph#:		
Allergist:		P	Phone #:		
Pediatrician/Family doctor:		Р	Phone #:		
Known allergies:					
Does child wear MedicAlert™ identification worn for asthma? ☐ YES ☐ NO					
TRIGGERS - List items that most commonly trigger your child's asthma.					
<u>RELIEVER MEDICATION</u> (or bronchodilators) provides fast temporary relief from asthma symptoms. It is recommended that reliever medication is carried with the child so it is available if an asthma episode occurs.					
What Reliever medication has been	☐ Salbutamo	Salbutamol (e.g. Ventolin [®] , Airomir [®])			
prescribed for your child? (CHECK ONE)	Symbicort	Oth	Other		
How many puffs of Reliever	1 puff	☐ 1 or	r 2 puffs		
medication_are prescribed for an asthma episode? (CHECK ONE)	2 puffs	othe	er		
Where does your child carry his/he		c Durs	se		
Reliever medication? (CHECK ONE	☐ backpack	othe	er		
Does your child know when to take their Reliever medication?		Can your child ta medication on th		ver Yes	
CIRCLE the type of medication device your child uses for Reliever medication.					
The second of th		(E-0)	PAGES 1		
Metered dose inhaler MDI & spa (MDI) with mouthp	niece MDI	& spacer Tu th mask	ırbuhaler [®]	Diskus [®]	

The Health Care Plan should accompany the child on excursions outside the facility.



ASTHMA HEALTH CARE PLAN

Birth date:		
S:		
 Remove child from triggers of asthma. Have child sit down. Ensure child takes Reliever medication (usually blue cap or bottom). Encourage slow deep breathing. Monitor child for improvement of asthma symptoms. If Reliever medication has been given and asthma symptoms do not improve in 5-10 minutes, contact parent/guardian or alternate contact. If any of the emergency situations occur (see list below), call 911/EMS. 		
 Activate 911/EMS. Delegate this task to another person. Do not leave the child alone. Continue to give Reliever medication every five minutes. Notify parent/guardian. Stay with child until EMS personnel arrives. 		
ey should inform the child's parent/guardian hal activities. ing. per week for asthma symptoms. polan on behalf of my child.		
nmunity program with required information. Date:		
n		

☐ Instruction sheet for medication device attached